



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{\omega \rho'}{1 + \sqrt{\rho'}} + \omega = \frac{S}{t} = \frac{\omega t'}{t}. \quad \therefore \rho' - \frac{t' - t}{t} \sqrt{\rho'} = \frac{t' - t}{t};$$

$$\therefore \sqrt{\rho'} = \frac{1}{2t} \left\{ (t' - t) + \sqrt{t'^2 - 3t^2 + 2tt'} \right\}.$$

$$\therefore \rho' = \frac{t' - t}{2t^2} \left\{ (t' + t) + \sqrt{t'^2 - 3t^2 + 2tt'} \right\} \dots \dots \dots (2).$$

Suppose Venus transits the sun's disc at such a rate that the sun's apparent diameter would be traversed in $7\frac{1}{2}$ hours, and at the same time the sun in his annual course moves through a distance equal to his own apparent diameter in 12 hours. Required (1) the distance from Venus to the sun, the earth's distance being unity, and (2) the distance from the earth to the sun, Venus's distance being unity.

Now $t = 7\frac{1}{2}$, $t' = 12$; hence for first case substituting in (1)

$$\rho = \frac{1}{4} \left(\frac{29}{4} - \sqrt{\frac{5}{4}} \right) = .721824.$$

For the second case substitute in (2)

$$\rho' = \frac{1}{4\frac{3}{4}} \{ 58 + \sqrt{1428} \} = 1.38538.$$

(The above is suggested in Proctor's Geometry of the Cycloid.)

A PROPOSITION IN DETERMINANTS.

By ALFRED HUME, C. E., D. Sc., Professor of Mathematics in the University of Mississippi.

THEOREM.—The product of two numbers, each the sum of four squares, is the sum of eight squares.

$$\begin{aligned} & \begin{vmatrix} a + b\sqrt{-1} & -c + d\sqrt{-1} \\ c + d\sqrt{-1} & a - b\sqrt{-1} \end{vmatrix} \times \begin{vmatrix} \alpha + \beta\sqrt{-1} & -\gamma + \delta\sqrt{-1} \\ \gamma + \delta\sqrt{-1} & \alpha - \beta\sqrt{-1} \end{vmatrix} \\ &= \begin{vmatrix} a + b\sqrt{-1} & -c + d\sqrt{-1} & 0 \\ c + d\sqrt{-1} & a - b\sqrt{-1} & 0 \\ 0 & 0 & 1 \end{vmatrix} \times (-1) \begin{vmatrix} \alpha + \beta\sqrt{-1} & 0 & -\gamma + \delta\sqrt{-1} \\ \gamma + \delta\sqrt{-1} & 0 & \alpha - \beta\sqrt{-1} \\ 0 & 1 & 0 \end{vmatrix} \\ &= (-1) \begin{vmatrix} a\alpha - b\beta + (a\beta + b\alpha)\sqrt{-1} & a\gamma - b\delta + (a\delta + b\gamma)\sqrt{-1} & -c + d\sqrt{-1} \\ c\alpha - d\beta + (c\beta + d\alpha)\sqrt{-1} & c\gamma - d\delta + (c\delta + d\gamma)\sqrt{-1} & a - b\sqrt{-1} \\ -\gamma + \delta\sqrt{-1} & \alpha - \beta\sqrt{-1} & 0 \end{vmatrix} \end{aligned}$$

$$= \begin{vmatrix} c\alpha - d\beta + (c\beta + d\alpha)\sqrt{-1} & c\gamma - d\delta + (c\delta + d\gamma)\sqrt{-1} \\ -c\gamma + d\delta + (c\delta + d\gamma)\sqrt{-1} & c\alpha - d\beta - (c\beta + d\alpha)\sqrt{-1} \end{vmatrix}$$

$$+ \begin{vmatrix} a\alpha - b\beta + (a\beta + b\alpha)\sqrt{-1} & a\gamma - b\delta + (a\delta + b\gamma)\sqrt{-1} \\ -a\gamma + b\delta + (a\delta + b\gamma)\sqrt{-1} & a\alpha - b\beta - (a\beta + b\alpha)\sqrt{-1} \end{vmatrix}$$

or $(a^2 + b^2 + c^2 + d^2)(\alpha^2 + \beta^2 + \gamma^2 + \delta^2) = (c\alpha - d\beta)^2 + (c\beta + d\alpha)^2$
 $+ (c\gamma - d\delta)^2 + (c\delta + d\gamma)^2 + (a\alpha - b\beta)^2 + (a\beta + b\alpha)^2 + (a\gamma - b\delta)^2 + (a\delta + b\gamma)^2.$

Euler's Theorem is an easy corollary of this, and *vice-versa*.

University of Mississippi, March, 1896.

A METHOD OF SOLVING QUADRATIC EQUATIONS.

By Prof. HENRY HEATON, M. Sc., Atlantic, Iowa.

Let it be required to solve the equation

$$ax^2 + bx + c = 0 \dots \dots \dots (1).$$

Transposing the middle term we have

$$ax^2 + c = -bx \dots \dots \dots (2).$$

$$\text{Squaring, } a^2x^4 + 2acx^2 + c^2 = b^2x^2 \dots \dots \dots (3).$$

$$\text{Subtracting } 4acx^2, \quad a^2x^4 - 2acx^2 = (b^2 - 4ac)x^2 \dots \dots \dots (4).$$

$$\text{Extracting the square root, } ax^2 - c = \pm (\sqrt{b^2 - 4ac})x \dots \dots \dots (5).$$

$$\text{Adding equation (2), } 2a^2x^2 = (-b \pm \sqrt{b^2 - 4ac})x \dots \dots \dots (6).$$

$$\text{Whence } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Let it be required to solve the equation $3x^2 - 2x = 21$.

Transposing $2x$ to the second member and 21 to the first, the equation becomes

$$3x^2 - 21 = 2x \dots \dots \dots (7).$$